










AcuPoint Analysis KIDNEY DISORDERS*

Points



AcuPoint Analysis
KIDNEY DISORDERS*
(DESCRIPTIONS)

Point	Description
C04	3 cun below, and in line with the navel.
GB25	At the waist on a "normal-waisted" person, along the sides of the body, nearly in line with the elbows on the lower border of your last rib.
K03	The width of two thumbs above the crown of the inner ankle, still on the inner side of the leg, but almost to the rear of the leg. Do not strongly stimulate this point after the third month of pregnancy.
K07	The width of two thumbs above and slightly behind the inside crown of the inner ankle.
SP06	The width of one hand (3 thumbs) above the crown of the inner ankle, just behind the shin bone on the front of the leg.
SP09	On the inner portion of the leg, just below the level of the kneecap, in the concave depression just above the calf, in line with the crown of the inner ankle.
UB23	1.5 cun from the spine on a level between the 2nd and 3rd lumbar vertebra level with the waist. Easily stimulated with the Back-eze.
UB28	1.5 Cun from the spine midline, between the lower medial border of the posterior superior iliac spine and the sacrum. 2 cun above the point where the buttocks form a crease. Easily stimulated with the Back-eze.
UB54	In the center of the crease at the rear of the knee, between the two ligaments. Do not use this point if you have varicose veins.