













AcuPoint Analysis MENSTRUATION ABSENCE OF

Points



AcuPoint Analysis

MENSTRUATION ABSENCE OF

(DESCRIPTIONS)

Point	Description
C04	3 cun below, and in line with the navel.
C06	1.5 Cun below the navel.
G01	In between the tip of the coccyx and the anus.
G04	On the spine atop the third lumbar vertebra, in line with the waist on a "normal-waisted" person.
K05	Between the crown of the inner ankle and the tip of the heel. The spot that is most pressure sensitive may be relatively large compared to most other points, and is found about 2 thumbs from the tip of the heel.
LI04	On the back of the hand between the thumb and index finger metacarpal bones. Find the spot midway on the bone that connects the index finger to the wrist. Sometimes this point is easy to find by squeezing two fingers together to form a small mound and marking an imaginary point at the peak of that mound. Then probe inward toward the main body of the hand, right on the bone.
LV08	On the inside of the knee, where the crease ends when the knee is bent.
SP06	The width of one hand (3 thumbs) above the crown of the inner ankle, just behind the shin bone on the front of the leg.
SP08	The width of one hand and two thumbs below the bottom of the kneecap, just behind the shin bone on the inner leg.
SP10	The width of two thumbs above the top of the kneecap, in the inner thigh, approximately in line with the crown of the inner ankle.
ST29	2 Cun from the median and 4 cun below the level of the navel.
ST36	The width of one hand (3 Cun) below the bottom of the kneecap, and the width of one thumb (1 Cun) toward the outside of the leg (direction of the small toe.) Found in the trough or valley where the two most prominent bones of the calf come together.
UB20	1.5 cun from the spine midline, between the 11th and 12th thoracic vertebra. 3.5 cun above the waist line on a 'normal-waisted' person. Easily stimulated with the Back-eze.
UB23	1.5 cun from the spine on a level between the 2nd and 3rd lumbar vertebra level with the waist. Easily stimulated with the Back-eze.
UB33	On the third posterior sacral foramen. 1 cun above the crease where the buttocks come together and 0.5 cun from the spine midline. A point easily stimulated with the Back-eze.
UB60	In the hollow (or valley) behind the crown of the outer ankle.